

Diverticulosis/Diverticulitis Diet

Diverticulosis: Characterized by the formation of pockets in the lining of the large intestine.



Diverticulitis: The acute stage of diverticulosis, when the pockets are infected or inflamed.

With the presence of diverticulosis it is best to follow a high fiber diet. This keeps pressure on the wall of the large intestine and prevents the accumulation of excess bacteria. Increase fiber gradually in your diet because sudden increases may cause increased formation of gas.


If diverticulitis does occur, after consulting with your physician, it may be recommended that you switch to a low fiber diet.



Recommended Foods for Diverticulosis

<u>Food Group</u>	<u>Amount to Consume Daily</u>	<u>High Fiber Foods</u>
Fresh Fruit 	2 or more servings	Apples (with skin), oranges, grapefruits, bananas, peaches, pears, plums, cherries, apricots and grapes. Avoid all berries with seeds, preserve with seeds, watermelon seeds and figs
Breads and cereals	4 or more servings	Whole grain bread (100% whole wheat, bran, or pumpernickel); dry or cooked cereals made from whole wheat, bran, oats, unprocessed wheat bran, or oat bran; whole grain crackers; bran muffins; brown rice. Avoid wheatberry bread, rye bread (with seeds), sesame, poppy, and sunflower seeds
Vegetables 	2 or more servings	Broccoli, brussel sprouts, peas, cabbage, carrots, celery, green beans, potatoes, zucchini, tomatoes (without seeds), onions, and baked beans. Avoid tomato seeds, corn, and popcorn
Miscellaneous	•	Avoid all nuts and chunky peanut butter

Recommended Foods for Diverticulitis

<u>Food Group</u>	<u>Amount to Consume Daily</u>	<u>Low Fiber Foods</u>
Fresh Fruit	2 or more servings	Fruit juice, fruit drinks, pureed fruit, cooked fruit (without skin)
Breads and cereals 	4 or more servings	White bread, French bread, farina, cream of wheat or rice, melba toast, saltines, corn flakes, rice krispies, puffed rice, white rice, pasta
Vegetables	2 or more servings	Vegetable juice, lettuce, winter squash, tomato sauce, mashed or cooked vegetables or potato
Miscellaneous	•	All dairy products, meat, poultry, fish, eggs, broth, cream soup

Diverticulosis & Diverticulitis

Not all people who have diverticulosis have symptoms or get diverticulitis

At least 50% of the population over 50 has diverticulosis



Symptoms can be mild and chronic or acute and severe

Diverticulosis

Pockets or pouches that form in the wall of the intestine

Symptoms

- Distention
- Bloating
- Spasms
- Cramping
- Constipation
- Pain may follow your meal
- Bowel movement may relieve pain
- Alternating diarrhea and constipation



Diverticulitis

Inflammation of the pockets and pouches caused by diverticulosis

Symptoms:

- Abdominal pain on the lower left side, frequently felt in the lower back
- Chills and Fever

Complications:

- Bleeding – a maroon or dark red blood may appear in your stool or in the toilet
- Obstruction or blockage in the large intestine, which may cause swelling
- Fistulas (connection of tissue between organs &/or skin) can cause infections
- Abscess – area that becomes infected with pus
- Peritonitis – abscess that ruptures and contents spill out into intestine

Provided as a service by
AstraZeneca

AstraZeneca 